

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005

arXiv:math/0504251v1 [math.GT] 12 Apr 2005



arXiv:math/0504251v1 [math.GT] 12 Apr 2005

2 -dimensional $\mathbb{Z}_n$ -manifold whose embedding $(F, F_s) \rightarrow (Z \setminus \partial Z, Z_s)$ induces an embedding of regular sets $F' \rightarrow Z'$. As M is closed and Z' is compact, F' has a finite number of connected components ([1], corollary 17.2(IV)).

In $Z = S^1 \times \text{Map}^n(S^1, S^1)$ consider the subspace $G = \{0\} \times \text{Map}^n(S^1, S^1)$ where $0 \in S^1$. We have $(G, G_s) \subset (Z, Z_s)$ and $G' \subset Z'$. So F' (which is union of surfaces with boundaries) and G' (which is an annulus) are embedded in $Z' = S^1 \times S^1 \times I$. Denote the boundary component of G' lying ∂Z by $\partial_0 G'$ and the other one by $\partial_1 G'$.




Figure 1: Regular parts, the covering $\partial_1 Z' \rightarrow Z_s$, and $Z_s \cup G \cup F$

Step 1: elimination of trivial circles of $\partial F'$. Denote the boundary component of Z' giving ∂Z by $\partial_0 Z'$ and the other one (participating in the identification) by $\partial_1 Z'$. Suppose that there is a circle of $\partial F'$ that is trivial in $\partial_1 Z'$. Then its projection on Z_s is trivial, too. As its projection is embedded in $Z_s \simeq T^2$ (because $\partial F'$ is the preimage of F_s which is embedded), it bounds an embedded disk there. Now we can proceed as in Proposition 1 of [4]: take a map of 2-disk in M bounding the same loop. As $\pi_2(W) = 0$, the union of these two disks bounds a map of 3-disk $\alpha : D^3 \rightarrow W$, which can be separated from M . Denote by N a small book neighborhood of $D^2 \subset Z_s$ in Z . The map f will not be changed on $\overline{Z} \setminus \overline{N}$; on D^2 the homotopy of f will be the pushing across $\alpha(D^3)$, and $N \setminus D$ will serve to rely the new map on D^2 and the old map on $\overline{Z} \setminus N$, doing it separately on each leaf of the book.




Figure 2: Changing of f on $D^2 \subset Z_s$.

This homotopy of f will eliminate the trivial circle from F' :




Figure 3: Surgery on a trivial loop of $\partial F'$.

Step 2: Reduction of F' to a union of closed surfaces and annuli. Now all the circles $\partial F' \subset \partial_1 Z' \simeq T^2$ are non-trivial and embedded, hence, parallel. They are either parallel to $\partial_1 G'$, or not, which corresponds respectively to $\partial F' \cap \partial_1 G' = \emptyset$ or $\partial F' \cap \partial_1 G' \neq \emptyset$. Take one connected component of F' , denote it still by F' .

We will separate two cases: $F' \cap G' = \emptyset$ or $F' \cap G' \neq \emptyset$.

Case 1: $F' \cap G' = \emptyset$. It implies $F' \subset \overline{(Z' \setminus G')} \simeq S^1 \times I \times I$. Note that $\overline{(Z' \setminus G')} \subset Z'$ is π_1 -injective but $F' \subset Z'$ is not, so neither is $F' \subset \overline{(Z' \setminus G')}$.

In this case $(F', \partial F') \rightarrow (S^1 \times I \times I, \partial(S^1 \times I \times I))$, $\partial F'$ going to a curve parallel to the generator of $\pi_1(S^1 \times I \times I) = H_1(S^1 \times I \times I)$. We need to find an embedded disk in $S^1 \times I \times I$ for the trivializing loop. As $\partial F'$ bounds $F' \subset S^1 \times I \times I$, $\partial F' \sim 0$ in $H^1(S^1 \times I \times I, \mathbb{Z}_2)$. As all the curves of $\partial F'$ are $\neq 0$ and parallel in $\partial(S^1 \times I \times I)$, they are in even number: say m with one orientation, m with the opposite one. $(F', \partial F') \rightarrow (S^1 \times I \times I, \partial(S^1 \times I \times I))$, so each $\partial F'$ is sent to a generator of $\pi_1(S^1 \times I \times I)$. Take an embedded disk $D = \{0\} \times I \times I \subset (S^1 \times I \times I)$. We have then $(D, \partial D) \rightarrow (S^1 \times I \times I, \partial(S^1 \times I \times I))$ and $D \cap F'$ is the union of circles and annuli. Take the innermost circle of $D \cap F'$; if it is ~ 0 in F' , we have a disk for the surgery, otherwise move D by isotopy to eliminate it, and so on. After treating all the circles either we have a disk for the surgery, or there are only arcs in $D \cap F'$. Take two components of ∂F corresponding to the arc whose ends are neighbouring in ∂D , denote them by a and a^{-1} . Then F' can be presented as $F' = A \cup_{\gamma} F''$ where A is an annulus with a hole, $\partial A = a \cup a^{-1} \cup \gamma$ and F'' is the remaining part.




Figure 4: F' as union of an annulus with a hole and F'' .

Denote by A' the annulus in $\partial(S^1 \times I \times I)$ lying between a and a^{-1} . Note that $\gamma \sim aa^{-1} \sim 0$ in $\pi_1(S^1 \times I \times I)$ (but $\gamma \neq 0$ in F' as $F' \neq S^1 \times I$). The arcs $A \cap D$ and $A' \cap D$ bound a disk in D , whose interior does not intersect F' .




Figure 5: $D \cap F'$ in D and D in $S^1 \times I \times I$.

Push $A \cup A'$ along its normal bundle toward the inside of this disk, living γ unchanged, and do the surgery of the pushed part (which is a torus with a hole) on the interior disk. We obtain a disk $\tilde{D}$ embedded in $S^1 \times I \times I$ such that $\partial \tilde{D} = \gamma$ and $\tilde{D} \cap F' = \gamma$. Hence we can move f by

of an annulus and $F \cap \partial D$. Doing this for every pair of components of $\partial F'$ corresponding to an arc with neighboring (in ∂D) ends, we will reduce F' to a union of closed surfaces (which will be treated as in the closed manifold case) and annuli.

Case 2: $F' \cap G' \neq \emptyset$ is a 1-dim manifold (embedded in both F' and G'); recall that G' was chosen such that $(F_s \cap G_s = \emptyset \Leftrightarrow \partial F' \cap \partial_1 G' = \emptyset)$.

Case 2.1: $F' \cap G' \neq \emptyset$ but $F' \cap \partial_1 G' = \emptyset$. Then $\partial F'$ is a union of circles parallel to $\partial_1 G'$; and $F' \cap G'$ is a closed 1-manifold. Circles of $F' \cap G'$ that are trivial in $\pi_1(F')$, can be eliminated by moving G' (without moving $\partial_1 G'$). So, if all the circles of $F' \cap G'$ are trivial in $\pi_1(F')$, we are back to the Case 1. If there is a circle of $F' \cap G'$ that is non-trivial in $\pi_1(F')$, it is either parallel to a component of $\partial F'$ or not.

- If there exists a circle not parallel to a component of $\partial F'$, then it is embedded in G' and not parallel to its boundary, so it bounds a disk in G' . We can do the surgery on this disk (choosing the innermost one).

- If all circles of $F' \cap G'$ are parallel to a component of $\partial F'$, then cutting Z' along G' , we'll have some number of π_1 -injective annuli embedded in $S^1 \times I \times I$, boundary going to boundary, hence isotopic to annuli in $\partial(S^1 \times I \times I)$ by an isotopy that is trivial on the boundary. Use these isotopies in Z' to move G' in order to separate it from F' , and we are again in the settings of the Case 1.

Case 2.2: $F' \cap G' \neq \emptyset$ and $\partial F' \cap \partial_1 G' \neq \emptyset$. If the components of $\partial F'$ are homotopic to $\partial_1 G'$, move G' by isotopy to separate $\partial_1 G'$ from $\partial F'$, and we are in the previous case. If the curves of $\partial F'$ are not homotopic to $\partial_1 G'$, choose an embedded curve $\alpha \subset \partial_1 Z'$, parallel to the curves $\partial F'$ (different from them) and make $\alpha \times I \subset Z'$ be the new G' . Then, we are back to Case 2.1.

Step 3: the image of the singular set Z_s can be separated from M . Now every connected component of F' is a torus or an annulus. Let us show that all the annuli can be eliminated. To simplify the notations, F will stand for F' without tori-components.

Fix a decomposition of $Map^n(S^1, S^1)$ by a wedge of intervals as follows:




Figure 6: Decomposition of $Map^n(S^1, S^1)$ on sheets.

After multiplying by S^1 , it gives a wedge of annuli $\bigvee S_i$ (identified along one boundary component) embedded in Z and decomposing Z into sheets $I \times I \times S^1$, in which the corresponding parts of F are manifolds. One generator of $\pi_1(Z_s)$ is given by the singular circle of $Map^n(S^1, S^1)$. The loop $(\bigvee S_i) \cap Z_s$ corresponds to the second generator, and can be chosen not to be parallel to F_s . Then every connected component of F intersects a decomposing annulus S_i by arcs. Take an arc coming from a component $F_1 \subset F'$, and suppose it is the innermost one in S_i (i.e. its union with an arc from ∂S_i bounds a disk in S_i that does not contain other arcs from $F \cap S_i$); denote it by $\alpha \subset F_1 \cap S_1$. Then the arcs $F_1 \cap S_j$ are the innermost ones in $S_j \forall j \neq i$. The product of I with the union of the corresponding disks in S_1 and S_2 (see Fig. 7 below)




Figure 7: Disks in the sheets.

allows to change f by homotopy so that the new image of the singular set Z_s coincide with the image of $I \times (F_1 \cap S_1)$. Hence by pushing off along the normal bundle of M , $\alpha \times I$ can be

S_1 and S_2 can be pushed away from M). Taking the product of the union of disks in S_k and S_{k+1} by I and doing the same homotopy for the sheet lying between S_k and S_{k+1} , one will eliminate all the annuli of F .

□

2 Application to 4-dim graph-manifolds

2.1 Seifert manifolds

Following [6, 7], we say that an orientable 4-manifold S with boundary is a *Seifert manifold* if it has the structure of a fibered orbifold $\pi : S \rightarrow B$ over a 2-orbifold with generic fiber T^2 , and S and ∂S are non-singular as orbifolds. Note that B as orbifold has no boundary, but the underlying surface of B does.

Local picture.

Any point $b \in B$ has a neighborhood of type $D = D^2/G$ where G is a finite subgroup of $\mathcal{O}(2)$ corresponding to the stabilizer of p . Then $\pi^{-1}(D) = (T^2 \times D^2)/G$ where the action of G is free and is a lifting of an action of G on D^2 (that is $\pi|_{\pi^{-1}(D)} : (T^2 \times D^2)/G \rightarrow D^2/G$ is induced by the canonical projection $T^2 \times D^2 \rightarrow D^2$).

Case 0: $G = \{1\}$.

In that case p is non-singular and $\pi^{-1}(p) = T^2$ is a regular fiber. The preimage of an arc in B that joins p with ∂B is $T^2 \times I$.


Case 1: $G = \mathbb{Z}_m = \langle g \mid g^m = 1 \rangle$ and the action is given by $g(x, y, z) = (x - a/m, y - b/m, ze^{\frac{2\pi i}{m}})$, where $(x, y) \in T^2$, $z \in D^2 = \{z \in \mathbb{C} \mid |z| \leq 1\}$, and m, a, b being mutually relatively prime. Then p is a cone point of angle $2\pi/m$ and $\pi^{-1}(p) = T^2$. The covering (regular fiber) $\rightarrow$ (singular fiber) corresponds to the subgroup $a\mathbb{Z} \oplus b\mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z}$. As $(a, b) = 1$, we can choose a basis in the π_1 of the regular fiber such that the covering corresponds to the subgroup $\mathbb{Z} \oplus n\mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z}$. Then the covering $T^2 \times D \rightarrow (T^2 \times D^2)/G$ splits as

$$\begin{array}{ccc} S^1 & \times & (S^1 \times D^2) \\ \text{id} \downarrow & & \downarrow \text{n-fold covering} \\ S^1 & \tilde{\times} & (S^1 \times D^2) \end{array}$$

(the bundle $S^1 \tilde{\times} (S^1 \times D^2)$ being trivial because $(T^2 \times D^2)/G$ is orientable). That means that $(T^2 \times D^2)/G = S^1 \times (3\text{-dimensional model})$. So in this case, the preimage of an arc in B that joins p with ∂B is the $\mathbb{Z}_n$ -manifold $S^1 \times \text{Map}^n(S^1, S^1)$.


Case 2: $G = \mathbb{Z}_2 = \langle \tau \mid \tau^2 = 1 \rangle$ and the action is given by $\tau \cdot (x, y, z) = (x + 1/2, -y, \bar{z})$. Then $\pi^{-1}(p) = K^2$, p lies on a reflector circle and $\pi^{-1}(D)$ is a twisted D^2 -bundle over K^2 . The preimage of an arc in B that joins the point p with ∂B is in this case a manifold $K^2 \tilde{\times} I$.

Case 3: $G = D_{2m} = \langle \tau, g \mid \tau^2 = g^m = 1, \tau g \tau^{-1} = g^{-1} \rangle$ and the action is given by $\tau \cdot (x, y, z) = (x + 1/2, -y, \bar{z})$ and $g \cdot (x, y, z) = (x, y - b/m, ze^{\frac{2\pi i}{m}})$. Then p is a corner reflector of angle π/m and $\pi^{-1}(p) = K^2$ which is m -fold covered by K^2 over a regular point of the reflector circle.



The preimage of an arc in B that joins p with ∂B is the mapping cylinder of a covering $T^2 \rightarrow K^2$ which is a 2-fold covering over the loop reversing the orientation of K^2 and a m -fold covering over the other loop. This is not a $\mathbb{Z}_k$ -manifold, but one can join p with ∂B by 2 consecutive arcs α and β , the first one lying on the reflector circle and not containing another corner reflector, the second one joining the other end of α with ∂B . Then $\pi^{-1}(\alpha) = S^1 \tilde{\times} \text{Map}^n(S^1, S^1)$ and $\pi^{-1}(\beta) = I \tilde{\times} K^2$.

As in dim 3, on the fundamental group level the singular fibers create roots of the regular fiber.



$S = M \cup M_S$, where M is a T^2 -bundle over a surface and M_S is its Seifert-part. Note that $0 \rightarrow \pi_1(M) \rightarrow \pi_1(S)$.

Theorem ([9], [8], [6], [7].) *Let S, S' be 4-dimensional closed orientable Seifert bundles with hyperbolic bases. Then $\pi_1(S) = \pi_1(S')$ if and only if there is a fiber-preserving diffeomorphism between S and S' .*

By taking the doubles, the previous theorem implies that if two Seifert manifolds with boundaries $(S, \partial S)$ and $(S', \partial S')$ over hyperbolic 2-orbifolds are homotopy equivalent rel boundary then S and S' are diffeomorphic.

2.2 4-dimensional graph-manifolds

A block is a Seifert bundle (with boundary) over a hyperbolic 2-orbifold. A graph-manifold structure on a compact closed oriented 4-manifold is a decomposition as a union of blocks, glued by diffeomorphisms of the boundary. Note that the boundary of a block has the structure of a T^2 -bundle over a circle. A graph-manifold structure is *reduced* if none of the glueing maps are isotopic to fiber-preserving maps of T^2 -bundles. Any graph-structure give rise to a reduced one by forming blocks glued by bundle maps into larger blocks. Like in the non-singular case, 4-dimensional graph-manifolds are aspherical, their Euler characteristic is 0 (because the blocks are finitely covered by T^2 -bundles over hyperbolic surfaces, hence have $\chi = 0$, and the glueings are made along 3-manifolds), and can be smoothed.

Theorem 1. *Any homotopy equivalence of closed oriented 4-manifolds with reduced graph-structures is homotopic to a diffeomorphism preserving the structures.*

We will say that a π_1 -injective map between the blocks $f : S \rightarrow S'$ is *fiber covering* if in the fundamental groups it sends the (normal) fiber subgroup of $\pi_1(S)$ into the (normal) fiber subgroup of $\pi_1(S')$. To extend the proof of the non-singular case ([4]), one has to show that any π_1 -injective map $f : W = \cup W_i \rightarrow W' = \cup W'_k$ of graph-manifolds with reduced graph-structures is homotopic to $\bigcup f_i$, where each $f_i : (W_i, \partial W_i) \rightarrow (W'_j, \partial W'_j)$ is a fiber covering map. The missing step is the following

Proposition 1. *Let S be a block and W be a 4-dim graph-manifold with reduced graph-structure. Then any π_1 -injective map $f : S \rightarrow W$ is homotopic to a map into one block of W .*

Proof. In the base of $\pi : S \rightarrow B$, take the wedge of circle on which B retracts. Join its core by arcs with every cone point and every reflector circle. As S retracts on

$$S' = \pi^{-1}(\text{wedge of circles} + \text{wedge of arcs} + \text{reflector circles}),$$

we have to show that $f|_{S'}$ is homotopic to a map into one bloc of W .

Step 1: $f|_{\pi^{-1}(\text{wedge of circles})}$ is homotopic to a map in one block of W . To make the reasonings as in the singular case, one only needs to show that up to homotopy rel boundary, any π_1 -injective map $g : (T^2 \times I, \partial(T^2 \times I)) \rightarrow (S, \partial S)$ is either fiber-covering or is a map into ∂S . Note that S is finitely covered by $\tilde{S}$ which is a T^2 -bundle over a hyperbolic surface (with boundary). As g sends boundary to boundary, $\text{Im } g_* \subset \pi_1(\tilde{S}) \subset \pi_1(S)$, hence g lifts to a map $\tilde{g} : (T^2 \times I, \partial) \rightarrow (\tilde{S}, \partial\tilde{S})$, to which Lemma 2 ([4]) applies.

Step 2: $f|_{\pi^{-1}(\text{wedge of arcs} + \text{reflector circles})}$ is homotopic to a map in one block of W . Denote by M the block of W in which $f(\pi^{-1}(\text{wedge of circles}))$ lies and denote p the point of B which is the common core of wedge of circles and wedge of arcs. Then $f|_{\pi^{-1}(p)}$ is a covering of the regular fiber of M .

Step 2.1: arcs to the cone points. The preimage of such an arc is $Z = S^1 \times \text{Map}^n(S^1, S^1)$, and $f|_Z$ sends ∂Z into M , hence $f(\partial Z)$ does not intersect the decomposing submanifolds of W .

and every component of $(f|_Z)^{-1}(\sqcup M_{\varphi_i})$ is a torus π_1 -injectively embedded in $Z = S^1 \times S^1 \times I$. Hence $(f|_Z)^{-1}(\sqcup M_{\varphi_i})$ is $\sqcup T^2$ that are parallel to ∂Z and hence are sent by f on the covering of regular fiber of M . Take the first T_1^2 from $\sqcup T^2$ counting from ∂Z , say it comes from $M_\varphi \subset M \cap M'$. If it exists, denote the next torus from $\sqcup T^2$ by T_2^2 . The restriction of f on $T^2 \times I \subset Z$ lying between T_1^2 and T_2^2 is a π_1 -injective map $f_1 : (T^2 \times I, \partial) \rightarrow (M', \partial M')$. As $f(T_i^2)$ is a covering of the regular fiber of M and the graph-structure is reduced, $f(T_i^2)$ is not a covering of the regular fiber of the neighbouring block M' , hence $f(T_2^2) \subset M_\varphi$ and f_1 is homotopic rel boundary to a map into M_φ . Continuing like this, we change $f|_Z$ so that $(f|_Z)^{-1}(\sqcup M_{\varphi_i})$ contains at most one manifold component. If there are none, we are done. If there is one such component, we have a π_1 -injective map $f : (Z, \partial Z, Z_s) \rightarrow (M \cup M', M, M')$.




Figure 8: $f(S^1 \times \text{Map}^n(S^1, S^1))$ in the blocs M and M'

As the torus $(f|_Z)^{-1}(\sqcup M_{\varphi_i})$ is parallel to ∂Z , we have a π_1 -injective map $f' : (Z, \partial Z) \rightarrow (M', \partial M')$, such that $f'(\partial Z)$ is not fiber-covering.

Case 1: f' lifts to a map into $\tilde{M}'$. If the torus $\tilde{f}'(\partial Z)$ corresponds to a $(\mathbb{Z} \oplus \mathbb{Z})$ -subgroup $\langle \alpha^n, \beta \rangle$ in $\pi_1(W)$, then $\tilde{f}'(Z_s)$ corresponds to the $(\mathbb{Z} \oplus \mathbb{Z})$ -subgroup $\langle \alpha, \beta \rangle$. Denote p' the projection of $\tilde{M}'$. If $p'_*(\alpha^n) = 0$ then $p'_*(\alpha) = 0$, too because $\pi_1(\tilde{B}')$ is free. Hence α is homotopic to a loop in the fiber of $\tilde{M}'$. If $p'_*(\alpha^n) \neq 0$ then $p'_*(\alpha) \neq 0$, and we can apply

Fact 1. *Let B be a surface with boundary, S a component of ∂B . Then the image $\pi_1(S) \rightarrow \pi_1(B)$ is root closed if and only if B is not a Möbius band and roots are squared.*

Proof. First note that by [3] roots are unique in free groups. If ∂B has more than one component, then S corresponds to a primitive element of $\pi_1(B)$, and the statement comes from [3] ($a^k = s^r$ implies that a, s are powers of a common element, and as a is primitive, s is a power of a).

Now suppose ∂B has one component. Denote $M = \text{Map}^k(S^1, S^1)$. Suppose there exists $a \in \pi_1(B)$ such that $a \notin \pi_1(\partial B)$ and $a^k \in \pi_1(\partial B)$. It gives a π_1 -injective map $(M, \partial M) \rightarrow (B, \partial B)$ such that $\pi_1(M) \rightarrow a$ and $\pi_1(\partial M) \rightarrow a^k$. Attach disks to M and B to get a π_1 -injective map $M \cup_{\partial M} D^2 \rightarrow B \cup_{\partial B} D^2$. As a^k is a finite power of a generator of $\pi_1(B)$, $\pi_1(M \cup_{\partial M} D^2) \rightarrow \pi_1(B \cup_{\partial B} D^2)$ is of finite index. As $\pi_1(M \cup_{\partial M} D^2) = \mathbb{Z}_k$ and $(B \cup_{\partial B} D^2)$ is closed, $(B \cup_{\partial B} D^2) \sim \mathbb{RP}^2$, because $\mathbb{RP}^2$ is the only closed surface with finite non-trivial π_1 . Hence $k = 2$ and B is the Möbius band. $\square$

Hence if $p'(\alpha^n)$ is homotopic to a loop in $\partial \tilde{B}'$, then so is $p'(\alpha)$. Hence α is homotopic to a loop in $\partial \tilde{M}'$, and f' is homotopic to a map into $\partial M'$.

Case 2: f' does not lift to a map into $\tilde{M}'$. Then α^n have roots on the singular fibers of M' , hence α^n is homotopic to a loop in the regular fiber of M' . Denote the Seifert projection of M' by π' . As the graph-structure is reduced and $\langle \alpha^n, \beta \rangle$ corresponds to the regular fiber of M , $\pi'_*(\langle \alpha^n, \beta \rangle) \neq 0$, hence $\pi'_*(\beta) \neq 0$. Which means that α can not commute with β because the roots of the regular fiber do not commute with the elements of the base.

Step 2.2: the image of a reflector circle without corner reflectors lies in one block.

The preimage of the reflector circle is twisted K^2 -bundle over S^1 . Applying the previous step to the preimage of the arc between ∂B and the reflector circle, we see that f restricted to the fiber of this bundle is homotopic to a map into the block M . Take in B a loop γ near the reflector

covers the regular fiber of the block M , hence the whole $f(M_{\pm id})$ can be shrunk by homotopy into M , in particular $f(\gamma)$ does. Hence the image of the whole 1-skeleton of $\pi^{-1}(\text{reflector circle})$ can be shrunk by homotopy into M , and the asphericity implies the statement.

Step 2.3 : case of reflector circles having corner reflectors. The preimage of an arc on a reflector circle between a corner reflector and a neighboring point is $\tilde{Z} = S^1 \tilde{\times} \text{Map}^n(S^1, S^1)$. Denote as before $\partial \tilde{Z} = S^1 \tilde{\times} \partial(\text{Map}^n(S^1, S^1))$ and $\tilde{Z}_s = S^1 \tilde{\times} (\text{Map}^n(S^1, S^1))_s$. The preimage of an arc in B between the reflector circle and the boundary is $I \tilde{\times} K^2$, and by previous its image by f can be shrunk into M .




Figure 9: Access to a corner reflector by two arcs.

Let us show that $f(\tilde{Z})$ such that $f(\partial \tilde{Z}) \subset M$ can be shrunk into M . The double covering $p : Z \rightarrow \tilde{Z}$ induces a π_1 -injective map $fp : Z \rightarrow W$ which by previous can be shrunk into M (by a homotopy which is constant on ∂Z). Let us show that the whole mapping cylinder of p , $\text{Map}_p(Z, \tilde{Z})$ can be shrunk into M . For this, note that the mapping cylinders of the 2-fold coverings $p|_{\partial Z}$ and $p|_{Z_s}$ are $(I \tilde{\times} K^2)$'s with boundaries lying in M , hence both can be shrunk into M . It remains to remark that the union of these mapping cylinders with Z contains a 1-skeleton of $\text{Map}_p(Z, \tilde{Z})$.

□

References

- [1] R. Abraham, J. Robbin, *Transversal mappings and flows*, New York-Amsterdam 1967.-161 p.
- [2] J. A. Hillman, *Four-manifolds, geometries and knots*, Geometry & Topology Monographs, 5. Geometry & Topology Publications, Coventry, 2002.-379 p.
- [3] R. C. Lyndon, M. P. Schützenberger, *The equation $a^M = b^N c^P$ in a free group*, Michigan Math. J. **9** (1962), pp. 289–298.
- [4] A. Mozgova, *Non-singular graph-manifolds of dimension 4*, available on <http://arXiv.org/abs/math.GT/0411335>
- [5] J. Stallings, *Group theory and three-dimensional manifolds*, A James K. Whittemore Lecture in Mathematics given at Yale University, 1969. Yale Mathematical Monographs, 4. Yale University Press, New Haven, Conn.-London, 1971.-65 p.
- [6] M. Ue, *Geometric 4-manifolds in the sense of Thurston and Seifert 4-manifolds. I*, J. Math. Soc. Japan **42** (1990), no. 3, pp. 511–540.
- [7] M. Ue, *Geometric 4 manifolds in the sense of Thurston and Seifert 4 manifolds. II*, J. Math. Soc. Japan **43** (1991), no. 1, pp. 149–183.
- [8] E. Vogt, *Vierdimensionale Seifertsche Faserräume*, Dissertation zur Erlangung des Grades eines Doktors der Naturwissenschaften der Abteilung für Mathematik an der Ruhr-Universität Bochum, Ruhr-Universität, Bochum, 1970. 151 p.
- [9] H. Zieschang, *On toric fiberings over surfaces*, Math. Notes **5** (1969), pp. 341–345.